AD)

Award Number: DAMD17-03-1-0157

TITLE: Dietary Determinants of Prostate Cancer

PRINCIPAL INVESTIGATOR: Lisa W. Chu, Ph.D.

CONTRACTING ORGANIZATION: The University of California

Berkeley, CA 94720

REPORT DATE: March 2006

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE	3. DATES COVERED (From - To)
01-03-2006	Annual Summary	1 Mar 200 3 – 28 Feb 2006
4. TITLE AND SUBTITLE		5a. CONTRACT NUMBER
Dietary Determinants of Prostate (Cancer	5b. GRANT NUMBER
		DAMD17-03-1-0157
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)	5d. PROJECT NUMBER	
Lisa W. Chu, Ph.D.		5e. TASK NUMBER
FM-" YWYGI OUT		55 WORK HAIT AHADED
E-Mail: <u>LWChu@lbl.gov</u>		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME	S(S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION REPORT
7.1 ERI ORIMINO OROANIZATION NAME	(c) AND ADDICEO(LO)	NUMBER
The University of California		
Irvine, California 94720		
9. SPONSORING / MONITORING AGENC		10. SPONSOR/MONITOR'S ACRONYM(S)
U.S. Army Medical Research and		7
Fort Detrick, Maryland 21702-501	2	
		11. SPONSOR/MONITOR'S REPORT
		NUMBER(S)
12. DISTRIBUTION / AVAILABILITY STA		
Approved for Public Release; Dist	ribution Unlimited	
40 OURDI EMENTARY NOTES		
13. SUPPLEMENTARY NOTES		
14. ABSTRACT		
14. ABSTRACT		
Prostate Canaar (PCa) access	unto for nearly 300/ of all nearly diama	sed cancers among American men. Epidemiologic
studies suggest that dietary fact	ore may be important in the stickers	f this disease. The objective of our research is to
determine how nutritional comp	ounds genistein betasitasteral (CIT) and	d omega-6 fatty acids (FA) function as modulators
accomme now munitional comp	Junus gematem, octasitosteror (511), all	d omega-o fatty acids (FA) function as modulators

Prostate Cancer (PCa) accounts for nearly 30% of all newly diagnosed cancers among American men. Epidemiologic studies suggest that dietary factors may be important in the etiology of this disease. The objective of our research is to determine how nutritional compounds genistein, betasitosterol (SIT), and omega-6 fatty acids (FA) function as modulators of PCa. In the third and final year of this fellowship, the fellow joined the National Cancer Institute's Cancer Prevention Fellowship Program and has continued working on the DOD-PCRP Postdoctoral Fellowship project with 30% effort (with knowledge and approval from the DOD-PCRP). During this year, we finished printing our custom microarray as well as optimized the hybridization protocol. We are currently performing expression analysis using the microarrays for both the *in vitro* and *in vivo* experiments using well established cell lines. Finally, we have initiated xenograft *in vivo* experiments with a novel PCa model that was developed recently in a collaborating laboratory.

1	5.	SU	JBJ	E	CT	Т	EI	RN	IS	

No subject terms provided.

16. SECURITY CI	LASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC					
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U	UU	8	19b. TELEPHONE NUMBER (include area code)					

Table of Contents

Cover1
SF 2982
Table of Contents3
Introduction4
Body4
Key Research Accomplishments8
Reportable Outcomes8
Conclusions8
Publicationsn/a
Appendicesn/a

Introduction

Prostate Cancer (PCa) accounts for nearly 30% of all newly diagnosed cancers among American men. Epidemiologic studies suggest that dietary factors may be important in the etiology of this disease. The objective of our research is to determine how nutritional compounds genistein, betasitosterol (SIT), and omega-6 fatty acids (FA) function as modulators of PCa. These three compounds belong to three distinct classes of dietary components, specifically isoflavonoid (genistein), phytosterol (SIT), and FA (omega-6 FA). Each class of compound could potentially modulate PCa in different ways and previous investigations have indicated that all three compounds may affect PCa in a dose-dependent fashion. We will determine differences in gene expression profiles at a range of doses for each compound on *in vitro* models. We will then test up to two doses on *in vivo* models to account for the complexity of tumor microenvironments.

In the first year, we began to develop the technical tools with which to investigate gene expression patterns that are modulated by the 3 dietary compounds. In the second year of study, we finished preparation of 355 cDNAs for fabricating the custom microarray for analysis of gene expression. In addition, we completed *in vitro* and *in vivo* experiments using well established PCa cell lines and the three dietary compounds in varying concentrations. In the third year, the fellow joined the National Cancer Institute's Cancer Prevention Fellowship Program and has continued working on the DOD-PCRP Postdoctoral Fellowship project with 30% effort (with knowledge and approval from the DOD-PCRP). During this past year, we finished printing our custom microarray as well as optimized the hybridization protocol. We are currently performing expression analysis using the microarrays for both the *in vitro* and *in vivo* experiments using well established cell lines. Finally, we have initiated xenograft *in vivo* experiments with a novel PCa model that was developed recently in a collaborating laboratory.

Here, we report our progress as it relates to the approved Statement of Work:

Task 1: To delineate the gene expression patterns of PCa in vitro of three nutritional compounds.

1.1 Obtain cDNAs for microarrays

From year 2, 344 cDNA clones from either in-house resources or ATCC have been prepared for our custom microarray. Genes chosen were tyrosine kinases or involved in cellular pathways with emphasis on androgen receptor (AR) and apoptosis pathways. Also included in our gene list are 3 control genes (GAPD, ACTB, and MBP) and other controls (Human Cot-1, Mouse Cot-1, Salmon Sperm DNA, and male human genomic DNA) for a total of 352 features.


1.2 Fabricate arrays and optimize protocols

We have successfully fabricated our custom microarray with the above mentioned 352 features (in duplicate) (Table 1). For hybridization, we evaluated incorporating Cyanine 3- (Cy3) and Cyanine 5-dUTPs (Cy5) (Amersham) in the control and test samples by 2 methods: direct incorporation during reverse transcription polymerase chain reaction (RT-PCR) and during random priming (RP) after RT-PCR. Hybridization using labeled cDNA resulted in weak signals in the Cy3 probes and little or no signals in the Cy5 probes. Hybridization using RP probes resulted in much stronger signals in both Cy3 and Cy5 channels. A representative example of an array hybridization is shown in Figure 1. For subsequent array experiments, RP probe labeling will be utilized after RT-PCR of total mRNA from each sample.

Table 1: Map of the custom microarray printing

_																											
	RAD	SB CDC	25A LMN	4 2	5 -	ETS 2	ACT A	RAF	ABL	TNFR	SPT	AN1	EPH EPH	A1	NTR NTR	KZ ERB	TLN	1 GZM	B XX	PPA PPA	FR -	- H. S	EPH 3	MsGe	nMale	IL2R	٥
3	2 P.S.	25 25	25A LMN	4 <u>q</u>	5 >	ETS 2	ACT A1	RAF	ABL	TNFR .	SPT	AN1	EPH BB	PA	NTR	Z 88	TLN	1 GZM	B XY	PPA o	8 분	- 蓝 5	EPH A3	a9sw	nMale	ILZR	2
,	D4G	ESR L	₽ P	15B	Ē	168	F G	CD4	- JOS	18A1 ESR	C10	RPA	GST	ΣĘ,	NR1	된 달	H H	2 3	a KR	<u> </u>	TRA TRA	= =	22 25	CAS	2	MP S	3
30	D40	ESR	ALOX	15B TRA	드	2 28	HPG D	CD4	, g	ESR	C10	RPA	GST	Z Ž	NR1	고 전:	표 %	3 중	aKR 5	<u> </u>	TRA CT	= =	C20	CAS	27	IMP SH	4
2	i į	CDC	2 KRT	18 PLA	H.	Z 7	YWH	BAG	PDG	TNFR	SF X Y	MER.	¥ .	T IN	SF6	₹ H 8	TNFR	P K	A A	ROS	MAD	i ii	ZAP	일 :	50	H H	
23	Ę	S CDC	2 KRT	18 PLA	3	<u>F</u> E	YWH	BAG	PDG	TNFR	NFK NFK	MER	≦ 5	TNFR	91 3	E E	DZ TNFR	SF X	A A F	ROS	MAD	7 <u>1</u> 11	ZAP	을 물	E .	HPR 8	
"	F S	USA	PPA PPA	N I	AR SE	15A	POV 1	MAP	置置	o §	F H	- 2	MAPK	FGF APPE	MYB	4	# H H	3 4 6	SRD 5A7	3 2	TNFS	E	5 6	₩ 8	3	DKFZ P434	
2	Z S	USA	PPA	됩	AR 5	2 2	POV 1	MAP	뮖	9	F H	- 5	MAPK	FGF	MYB	3 3	# # E	3 4 6	SRD 547	IBG.	TNFS	E	6 6	器 5	3	DKFZ P434	
20	188	¥ 8	g 8	GAP GAP	۵	SERP	TP5 3	SF10	Ä	CAS P7	MAP	ξ,	4 型 8	3 # E	CAS	E K S	SINFS	DFF PFF	a L	Z Z	ANX A5	¥	EPH A	HuGe	nwaie	KRA S3	CCN T1
	E	¥.8	3 8	GAP	۵	NE1	TP5 3	SF10	Ě	CAS P7	MAP	ξ,	4 量 8	S	CAS	E K	TNFS	H H	٦ E	EM S	ANX A5	Ĕ	EPH A4	HuGe	an and an and an	KRA S2	CCN T1
18	CAS P4	5	BCA 55	CAS	P2 TNFR	8F10	TAB 1	ESR 1	ITGA 5	, ag	ALO X	등 -	- AF	H H H	BCL 241	NOT		3 8 5	RN 4	TGM 4	TNFR SF21	PPA P2B	PPA P2A	HH 5	3	TMPR	85 85
4	CAS P4		BCA E	CAS RS	P2 TNFR	B B	TAB 1	ESR 1	ITGA 5	RGR A	ALO x5	- GH	NFK	문 문 문	BCL 2A1	NGH		3 8 2	₩ 4	TGM 4	TNFR SF21	PPA P28	PPA P2A	뚦 5	3	TMPR SS2	P8 KB
16	M2 M2	SEX	§ 5		MBP	PA12	TNF SF6	DFF A	FGF R2	E E	MAP 3K14	YTY ^	Ę	E E	H 2	MET	TRA 2	TNFR SF10	S ∞	KDR	FLT3	TGB 3	ROR 2	CDK N2A		IGF1	MAD H2
15	MD M2	SDK	ğ ç	٥	MBP	PA12	TNF SF6	DFF A	FGF R2	ER	MAP 3K14	AT ~	Ě	Ä	HH 75	MFT	TRA 5:3	SF10	S R	XOX.	FLT3	1TGB 3	ROR 2	CDK N2A		GF1	MAD H2
14	DFF A	BCL 7.1	IL2R	CAS	CAS P9	F	G9 G9	DED D	MIC	HGF	BMX	ANG PT1	MTA	HNF 38	BCL 2	HEVI	BCL 211	CRA GG	TUB A3	AR	TNFR SF10 B	BAX	PSC A	¥		왕 9	LDH A
13	DFF A	집	L2R □	CAS	CAS PB	2	G9	DED	MIC	HGF	BMX	ANG PT1	MTA	AN AS	BCL 2	HEVI	BCL 2111	8 G	A3 TUB	AR	SF10 B	BAX	PSC A	¥		SHS G	LDH
12	CDC 25A	TP5	DIAB	S. S.	SZ5	KAIT	SF10 O	FGF 7	R FR	IGF1	KBK	KIT	RYK	EPH A7	CAS P10	SRM	Ľ	BCL ZA1	ACK	CSF 1R	ESR RG	N A	DKFZ P434 C141 8	PLA B	SSGe	N33	PDG FA
Ŧ	CDC 25A	TP5	DIAB	PRS PRS	225	KA!	SF10 D	FGF 7	RIK3	IGF1	ÄΒ. Θ	Ä	RYK	EPH A7	CAS P10	SRM	H	BCL 2A1	ACK	CSF 1R	ESR RG	N A	DKFZ p434 C141 8	PLA B	SSGe	N33	PDG
9	TNFS F12	EIF2 S1	SOS	¥.	₹ 1	8	SRF	CTN NB1	ITGA V	VEG	ALO X12	ANG PT2	CDC 37	AAT	ATM	0DC	KCN MB1	CDK N18	MDH 1	CSN K2B	SF10 A	BAK 1	BCR P1	BID	CAS P9	SRE BF1	ACP P
6	TNFS F12	EIF2 S1	SOS	¥ 5	₹ 1	8	SRF	CTN NB1	TGA V	VEG F	ALO X12	ANG PT2	CDC 37	AAT	ATM	0DC	KCN MB1	CDK N1B	MDH 1	CSN K2B	SF10 A	PAK	BCR P1	BID	CAS P9	SRE BF1	ACP P
	BRC A1	SDK N18	CDK N2A	F.	KISS	-	SFN	cyc s	MAP 2K1	SRC	ALO X12	ITGA 6	ТХК	ABL 2	CAS P7	MST 1R	YES 1	TNFR	NTR KR1	BTK	GB	IIGA 2B	XX.	주 참	SSGe	MYB	HNF 3G
~	BRC A1	SDK	NZA CD	F.F.	KISS	-	SFN	s CAC	MAP 2K1	SRC	ALO X12	ITGA 6	ТХК	ABL 2	CAS P7	MST 1R	YES 1	TNFR	MTR KR1	ВТК	읆	ZB ZB	FR	참 참	SSGe	MYB	HNF 3G
9	4KT	FAD	ALOX 5AP	88 -	GAS	2	WI/	MAD H4	ICA M1	EGF	NUM A1	BIR C5	CCN Pd	STA T1	ADP RT	SLK 3	ADA M9	CAS P3	NS 4	GAG EB1	¥.	ARH GDIB	SFR P4	ANX A7	CAS P8	ADM	불호
2	- AKT	FAD	ALOX 5AP	EBB -	GAS	7	MIN	MAD H4	M CA	EGF	NUM A1	器 &	CCN Pd	STA T1	ADP RT	KLK 3	ADA M9	CAS P3	NCO A4	GAG EB1	Ą	GDIB	SFR P4	ANX A7	CAS P8	ADM	불호
4	KLK 2	<u>Ş</u> 4	CDK N1A	PDP K1	NME	4	KAS A1	P4	S 8	ERB B2	ABC C6	YWH	R3 R3	ACT B	S. C.	EPH 81	PTK 9	PAR P	A Z	A2	9 MMP	FIT	SYK	ALK	MsC ot1	PRK CB1	S S
8	KLK2	Q 4	CDK N1A	PDP	NME	4 0	KAS PA	F 5	S S	ERB B2	ABC	YWH	FGF R3	ACT B	8 G	EPH B1	PTK 9	PAR P	문 도	A2 A2	9 B	FTT	SYK	ALK	MsC ot1	RK CB1	S S
2	- MMP	P8 SF	MXI1	s HR	90	o E	1.6A	2 E	ITGA 3	THB S2	BAD	ALD HI	PCN A	SERP	CSK	TA A	85 es	CAS P10	K10	물 4	PMA IP1	WE H	FAT	5A B	CAS P6	CST	TGF B111
-	M F	CAF P6	MX -	HRA S	DED	Ę	5 A 5	BP3	A3	THB S2	BAD	ALD H1	PCN A	SERP	CSK	RAP 14	PS FKB	CAS P10	MAP K10	g 4	PMA PT WX	HAE	FAT	54 A2	CAS P6	CST B	TGF B111
	-	2	3	4		,	9	7	80	6	10	Ξ	12	13	4	15	16	17	- 8	19	20	21	22	23	24	52	26

Figure 1: Representative hybridization patterns of RP labeled cDNAs in Cy3 channel (A), Cy5 channel (B) and Cy5/Cy3 ratio (C).



1.3 Grow cells under test conditions, isolate RNA, RT-PCR, label probes

We completed the cell culture experiments of this task in year 2. However, we were unable to replicated experiments using the test compound SIT due to difficulties in consistently dissolving SIT into solution with its appropriate vehicle (Cyclodextrin; Sigma). Because we successfully tested this compound in our *in vivo* xenograft model pilot study (Task 2), we have decided to delay the *in vitro* experiments using SIT until we are able have appropriate resources to reliably perform the experiments.

1.4 Perform microarray experiments, analyze data with standard statistical programs and Perform multicolor spectral transcript analysis and analyze data with standard statistical programs

Due to the time needed to optimize the hybridization protocol, we are currently in the process of performing array experiments and analyzing the expression patterns from RNA isolated from the *in vitro* experiments. See Figure 2 under Task 2 for examples of array hybridizations. The multicolor spectral transcript analysis will need to be delayed until all array experiments have been completed.

Task 2: To determine if and how three select dietary components modulate the growth and gene expression of clinical PCa specimens in vivo.


2.1 Pilot study to investigate timing of nutrient supplementation in SCID mice
In parallel with the *in vitro* experiments mentioned in Task 1, we performed *in vivo* experiments using xenograft technique on SCID mice at the comparator of the com

experiments using xenograft technique on SCID mice at the co-mentor's laboratory using the same cell lines mentioned above in year 2.

2.2 Analyze tumors from pilot study

Similar to task 1.4, we are currently performing array experiments and analyzing the expression patterns from RNAs isolated from the *in vivo* experiments. Examples of hybridizations using RNAs from this pilot study are shown in Figure 2. Likewise, the multicolor spectral transcript analysis will need to be delayed until all array experiments have been completed.

Figure 2: Representative hybridization patterns of CAFTD04 derived tumors grown in SCID mice fed with control diet (A) and Genistein supplemented diet (B). All test probes are labeled in Cy3 (green) and reference probes (generated from CAFTD04 derived tumors grown in SCID mice fed with control diet) are labeled in Cy5 (red).



2.3 Grow 6 tumors in 21 SCID mice (2 tumors per mice) that ingest one of 7 test diets, harvest tissues, prepare tissue sections and RNAs, RT-PCR, and make probes. Test diets include control and each compound at one of 2 concentration

For this task, we found it difficult to obtain clinical samples that were large enough to be implanted into the number of mice that was initially proposed. To circumvent this problem, we decided to use a newly generated PCa model that was developed by a collaborator (Wang et al., submitted) as mentioned in the previous report. Briefly, pieces of PCa tissue from a patient were grafted into the subrenal capsule site of testosterone-supplemented male SCID mice. After 5 serial transplantations, the tissues were transferred into mouse prostates. A metastatic tumor line generated from lymph nodes, designated PCa1-met, had few chromosomal alterations, as indicated by Spectral Karyotyping. Orthotopic grafting of PCa1-met in 47 hosts led in all cases to metastases to multiple organs (lymph nodes, lung, liver, kidney, spleen and, notably, bone). Histopathological analysis showed strong similarity between orthotopic grafts and their metastases which were of human origin as indicated by immunostaining using antibodies against human mitochondria, androgen receptor, prostate-specific antigen, pan-cytokeratin, p63, cytokeratin-8 and Ki-67. This model is unique due to the fact that the xenograft implants metastasized to regions of the mouse that paralleled the locations of clinical PCa metastases (most notably lung, liver and bone) in humans. In addition, the resulting cell lines contain few cytogenetic aberrations, similar to clinical PCa and unlike most well established cell lines (i.e., PC3, LNCaP, and DU145).

When we implemented the experiments using this new model, however, the SCID mice were adversely affected in unanticipated ways. At the same time, our collaborators were having similar problems. All animal experiments were stopped so that the issue can be resolved without harm to any more animals. Recent experiments as communicated by our collaborators showed that the murine stroma, which was contained within the tumors that are regrafted, harbored a mouse virus that infected the host SCID mice. The contaminated tumor lines have since been discarded and several un-infected tumor grafts have been established. We are currently in the process of using the new and un-infected tumor lines for the purpose of this study.

2.4 Perform microarray experiments, analyze data with standard statistical programs

Due to the problem described above, we are currently in the process of performing the in vivo experiments using the newly established PCa model. Once these experiments are complete, we will analyze the expression profiles using our custom arrays.

Key Research Accomplishments for Year 2

- Fabricated a custom microarray with 352 features in duplicate.
- Optimized microarray hybridization conditions.
- Began microarray expression profiling of RNAs from *in vitro* experiments with established PCa cell lines grown in control and test media containing different concentrations of test nutritional supplements.
- Began microarray expression profiling of RNAs from *in vivo* experiments in the comentor's laboratory with xenografts of well established PCa cell lines in SCID mice whose diets are supplemented with various test compounds.
- Initiated *in vivo* experiments in co-mentor's laboratory with xenografts of the newly developed and un-infected PCa tumor lines in SCID mice whose diets are supplemented with various test compounds.

Reportable Outcomes

- Became a National Cancer Institute Cancer Prevention Fellow (2005-2008)
 - O Beginning in July 2005, the fellow began a 1 year academic program to obtain a Masters in Public Health with an emphasis on Epidemiology at the University of California, Berkeley to supplement the fellow's training in the field of cancer prevention. Degree is expected to be conferred in May 2006.
 - Beginning in June 2006, the fellow will begin a 2 year research program at the National Cancer Institute in Bethesda, MD, which will focus on cancer prevention.

Conclusion

We have used the third year of this fellowship to fabricate our custom microarray with 352 features (in duplicate) for expression profiling of cells grown *in vitro* and *in vivo* under different dietary supplement conditions including none, genistein, SIT, and omega-6 FA. In addition, this year has been used to troubleshoot many unexpected issues such as microarray probe generation as well as viral infection of the new PCa tumor line models. Both major issues have been resolved. Microarray analysis for task 1 and the pilot study for task 2 are underway. Additionally, the *in vivo* xenograft model using the new PCa tumor lines have been initiated and expression profiling of the resulting tumors will be performed once tumors are harvested from co-mentor's laboratory.

Publications

None: pending final data analyses.